

## Moderna 3<sup>rd</sup> doses – Third dose is not the same as a Booster Dose

Third doses are available for those who are moderately and severely immunocompromised. Third dose can be administered 28 days (or later) after the 2nd dose. Conditions and treatments include but are not limited to:

- Active treatment for solid tumor and hematologic malignancies
- Receipt of solid-organ transplant and taking immunosuppressive therapy
- Receipt of chimeric antigen receptor (CAR)-T-cell or hematopoietic stem cell
- Moderate or severe primary immunodeficiency
- Advanced or untreated HIV infection
- Active treatment with high-dose corticosteroids

## Booster Doses – Moderna

For individuals who received a Moderna COVID-19 vaccine, the following groups are eligible for a booster shot at **6 months** or more after their initial series:

- 65 years and older
- Age 18+ who live in long-term care settings
- Age 18+ who have the following underlying medical conditions:
  - Cancer
  - Chronic kidney disease
  - Chronic liver disease
  - Chronic lung diseases
  - Dementia or other neurological conditions
  - Diabetes
  - Down syndrome
  - Heart Conditions
  - HIV infection
  - Weakened immune system
  - Mental health conditions
  - Overweight and obesity
  - Pregnancy
  - Sickle cell disease or thalassemia
  - Smoking (current or former)
  - Solid organ or blood stem cell transplant
  - Stroke or cerebrovascular disease
  - Substance use disorder,
  - Tuberculosis (TB)
- Age 18+ who work or live in high-risk settings such as:
  - First responders (e.g., healthcare workers, firefighters, police, congregate care staff)
  - Education staff (e.g., teachers, support staff, daycare workers)
  - Food and agriculture workers
  - Manufacturing workers
  - Corrections workers
  - U.S. Postal Service workers
  - Public transit workers
  - Grocery store workers

## Booster Doses – Pfizer

Anyone who previously received the Pfizer vaccine AND who falls in one of the categories below can receive a booster dose, as long as it has been at least six months since the second dose:

- Anyone 65 years old and older
- Anyone who lives in a long-term care facility who is at least 18 years old
- Anyone age **18-64** who has one of the following conditions:
  - Overweight or obesity
  - Diabetes
  - Long term (chronic) Kidney disease
  - High blood pressure
  - Long term lung disease like COPD
  - Dementia
  - Down's syndrome
  - Heart conditions
  - HIV
  - A weakened immune system
  - Liver disease
  - Pregnancy
  - Sickle cell disease
  - History of organ transplant
  - History of smoking
  - Stroke
  - Substance use disorder
- Anyone age **18-64** who is at high risk for COVID-19 exposure due to their job as a:
  - First responder (healthcare workers, firefighters, police, congregate care staff)
  - Education staff (teachers, support staff, daycare worker)
  - Food and agriculture worker
  - Manufacturing worker
  - Corrections worker
  - U.S. Postal Service worker
  - Public transit worker
  - Grocery store worker

## Booster Dose – J&J

- Anyone 18+
- At least 2 months since primary dose